



Please check the examination details below before entering your candidate information

Candidate surname

Other names

**Pearson Edexcel
Level 3 GCE**

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--

Monday 8 June 2020

Afternoon (Time: 1 hour 45 minutes)

Paper Reference **9CH0/02**

Chemistry

Advanced

Paper 2: Advanced Organic and Physical Chemistry

**Candidates must have: Scientific Calculator
Data Booklet
Ruler**

Total Marks

Instructions

- Use **black** ink or **black** ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
- *there may be more space than you need.*

Information

- The total mark for this paper is 90.
- The marks for **each** question are shown in brackets
- *use this as a guide as to how much time to spend on each question.*
- For the question marked with an **asterisk (*)**, marks will be awarded for your ability to structure your answer logically showing the points that you make are related or follow on from each other where appropriate.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Check your answers if you have time at the end.

Turn over ►

P62669A

©2020 Pearson Education Ltd.

1/1/1/1/1/



www.ibworldwideacademy.com



Pearson



Answer ALL questions.

**Some questions must be answered with a cross in a box ☒.
If you change your mind about an answer, put a line through the box ☒
and then mark your new answer with a cross ☒.**

1 This question is about methanol, CH_3OH .

(a) Draw a dot-and-cross diagram to show the bonding in a molecule of methanol.
Show outer shell electrons only.

(2)

(b) Predict which bond has the shortest bond length in a molecule of methanol.

(1)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(c) Methanol is soluble in water.

(i) State the strongest type of intermolecular force that occurs between molecules of methanol and water.

(1)

(ii) Draw a labelled diagram to show the interaction named in (c)(i) between one molecule of methanol and one molecule of water. Include any relevant lone pairs and dipoles in your diagram.

(3)

(Total for Question 1 = 7 marks)



P 6 2 6 6 9 A 0 3 3 2



2 This question is about alkenes.

(a) Which of these has the molecular formula C_6H_{10} ?

(1)


A



B




C



D



(b) What is the systematic name of this alkene?



(1)

A 2-methylpent-1-ene

B 3-methylpent-1-ene

C 2,3-dimethylbut-1-ene

D 2,3-dimethylbut-3-ene

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA



P 6 2 6 6 9 A 0 3 3 2



(c) Two reactions of ethene are shown.



Complete the table.

(3)

Reaction	Reagent and condition	Product
1	HBr at room temperature	
2		$\begin{array}{c} \text{H} & \text{H} \\ & \\ \text{H}-\text{C}-\text{C}-\text{OH} \\ & \\ \text{H} & \text{H} \end{array}$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 6 2 6 6 9 A 0 3 3 2




DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(d) But-1-ene has the structure



(i) Draw the structure of the polymer formed when but-1-ene polymerises. Include **two** repeat units.

(1)

(ii) Calculate the number of molecules in 70.0 g of but-1-ene.
[Avogadro constant = $6.02 \times 10^{23} \text{ mol}^{-1}$]

(2)

(Total for Question 2 = 8 marks)



P 6 2 6 6 9 A 0 3 3 2



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE




P 6 2 6 6 9 A 0 3 3 2



3 This question is about the compound potassium bromate, KBrO_3 .

(a) These bromate ions react with bromide ions in acidic solution.



(i) Explain, in terms of oxidation numbers, whether or not this is a disproportionation reaction.

(2)

.....

.....

.....

.....

.....

.....

(ii) What is the overall order of this reaction?

(1)

- A 3
- B 6
- C 12
- D cannot tell from this information

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA



P 6 2 6 6 9 A 0 3 3 2



(b) Potassium bromate decomposes on heating.



Calculate the maximum volume of oxygen, in dm^3 , measured at room temperature and pressure (r.t.p.), that could be produced from the complete decomposition of 5.20 g of potassium bromate.

[Molar volume of gas at r.t.p. = $24.0 \text{ dm}^3 \text{ mol}^{-1}$]

(3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 3 = 6 marks)




P 6 2 6 6 9 A 0 3 3 2



4 This question is about the identification of some organic compounds.

(a) The skeletal formulae of four organic compounds are shown.



(i) Which of these compounds can be hydrolysed to form methanol as one of the products?

(1)

- A** Compound **P**
- B** Compound **Q**
- C** Compound **R**
- D** Compound **S**

(ii) Which of these compounds produces carbon dioxide when it reacts with aqueous sodium hydrogencarbonate?

(1)

- A** Compound **P**
- B** Compound **Q**
- C** Compound **R**
- D** Compound **S**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 6 2 6 6 9 A 0 1 1 3 2



(b) Compound **T**, $C_4H_{10}O$, is oxidised by acidified potassium dichromate(VI) to form compound **U**, C_4H_8O .

U gives an orange precipitate with 2,4-dinitrophenylhydrazine (Brady's reagent) but does **not** give a red precipitate when heated with Fehling's solution.

T reacts with ethanoyl chloride to form compound **V**, $C_6H_{12}O_2$.

Deduce the structures of compounds **T**, **U** and **V**. Justify your answers.

(6)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(Total for Question 4 = 8 marks)





5 This question is about hydrocarbons.

(a) Which of these molecular formulae represents a non-cyclic, saturated hydrocarbon?

(1)

- A C_6H_6
- B C_6H_{10}
- C C_6H_{12}
- D C_6H_{14}

(b) How many **structural** isomers are there with the molecular formula C_5H_{12} ?

(1)

- A 2
- B 3
- C 4
- D 5

(c) How many σ bonds and π bonds are there in one molecule of cyclohexene?



(1)

	σ bonds	π bonds
<input type="checkbox"/> A	5	2
<input type="checkbox"/> B	6	1
<input type="checkbox"/> C	15	2
<input type="checkbox"/> D	16	1

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 6 2 6 6 9 A 0 1 1 3 2



(d) When hydrocarbons undergo complete combustion, there is a change in the total volume of gases.

(i) Ethane burns in excess oxygen.



All gas volumes are measured at the same temperature and pressure when water is a gas.

What is the **increase** in the total volume when 100 cm^3 of ethane is burned in excess oxygen?

(1)

- A 50 cm^3
- B 100 cm^3
- C 200 cm^3
- D 500 cm^3

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA




P 6 2 6 6 9 A 0 1 1 3 2



- (ii) A combustion experiment was carried out using conditions under which water was a liquid.

A cyclic hydrocarbon, C_xH_y , was mixed with excess oxygen and ignited. Under the conditions of the experiment, this hydrocarbon was gaseous and had a volume of 25 cm^3 .

The equation for the complete combustion of C_xH_y is



The total gas volume **decreased** by 75 cm^3 .

The remaining gases were shaken with aqueous sodium hydroxide and the total gas volume **decreased** by a further 125 cm^3 .

All gas volumes were measured at the same temperature and pressure.

Suggest the identity of the cyclic hydrocarbon by calculating the molecular formula of C_xH_y .

Include the **skeletal formula** of the cyclic hydrocarbon.

(3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





(e) Propene reacts with iodine monochloride, ICl, by an electrophilic addition mechanism.

Draw the mechanism for the reaction between propene and iodine monochloride to form the **major** product.

Include the dipole on the ICl molecule, curly arrows and any relevant lone pairs of electrons.

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 6 2 6 6 9 A 0 1 1 3 2



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(f) Limonene is obtained from the oil in lemon peel and it is the only alkene present.



0.500 g of the oil reacted with exactly 30.6 cm³ of a solution of bromine dissolved in cyclohexane with a concentration of 0.200 mol dm⁻³.

Calculate the percentage by mass of limonene in the oil.
Give your answer to an appropriate number of significant figures.

Assume that there is nothing else in the oil that reacts with bromine.

(4)

(Total for Question 5 = 15 marks)



P 6 2 6 6 9 A 0 1 1 3 2



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA


BLANK PAGE



P 6 2 6 6 9 A 0 1 1 3 2



- 6 A bromoalkane, RBr, reacts with aqueous hydroxide ions in a nucleophilic substitution reaction.



This reaction is first order with respect to the bromoalkane and the rate equation is

$$\text{rate} = k[\text{RBr}]^1[\text{OH}^-]^x$$

where x is the order of the reaction with respect to hydroxide ions.

In an experiment, a sample of the bromoalkane was added to a large excess of aqueous sodium hydroxide and the concentration of the bromoalkane was determined at regular time intervals.

Results

Time / s	[RBr] / mol dm ⁻³
0	0.100
30	0.065
60	0.042
90	0.028
120	0.019
150	0.014

- (a) This experiment is carried out using the bromoalkane dissolved in ethanol and the hydroxide ions dissolved in water.

Give a reason why a solution of hydroxide ions dissolved in pure ethanol should **not** be used.

(1)

.....

.....

.....

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA





(b) Plot a graph of $[RBr]$ against time.

(3)



(c) Explain how the graph shows that the reaction is first order with respect to RBr . Include the values of two consecutive half-lives.

You **must** show your working for the half-lives on the graph.

(2)

.....

.....

.....

.....

.....

.....

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 6 2 6 6 9 A 0 1 1 3 2




DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

- (d) The experiment was repeated using equal concentrations of RBr and varying the concentration of hydroxide ions.

A graph was plotted of the results.



- (i) Deduce the value of x in the rate equation.

$$\text{rate} = k[\text{RBr}]^1[\text{OH}^-]^x \quad (1)$$

- (ii) Give the mechanism for the reaction that is consistent with the orders of reaction with respect to R-Br and hydroxide ions. Include curly arrows and relevant lone pairs.

(3)





(e) 2-bromobutane can react with aqueous hydroxide ions by an S_N1 mechanism.

Explain why the butan-2-ol produced from a single optical isomer of 2-bromobutane, using this mechanism, is **not** optically active.

(3)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(Total for Question 6 = 13 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA




P 6 2 6 6 9 A 0 1 1 3 2



7 This question is about the synthesis of organic compounds.

(a) A student suggested the following plan for the synthesis of pentanoic acid from but-2-en-1-ol.



(i) LiAlH_4 is a source of hydride ions, H^- .

Give a possible reason why LiAlH_4 cannot be used to reduce alkenes.

(1)

.....

.....

.....

(ii) Give a suitable reagent and condition for Step 1.

(2)

.....

(iii) Step 2 is incorrect because alcohols can only be converted to nitriles via an intermediate compound.

Identify a suitable intermediate compound by name or formula.

(1)

.....

(iv) Step 3 involves the hydrolysis of a nitrile.

Give the additional reagent that should be added after heating under reflux with aqueous sodium hydroxide, to produce pentanoic acid.

(1)

.....

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





- (b) Devise a four-step synthesis, involving the use of a Grignard reagent, to convert benzene into benzoyl chloride.



Include the reagents and conditions for each step in the synthesis and the structures of the intermediates.

(7)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 7 = 12 marks)



P 6 2 6 6 9 A 0 1 1 3 2



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



P 6 2 6 6 9 A 0 1 1 3 2



8 This question is about the analysis of organic compounds.

(a) **X** is an organic compound.

(i) The accurate relative atomic masses, A_r , of the four elements that could make up **X** are shown in the table.

Element	A_r
hydrogen, H	1.0078
carbon, C	12.0000
nitrogen, N	14.0031
oxygen, O	15.9949

X gives a molecular ion peak at $m/z = 100.0522$ on its mass spectrum.

Which is the molecular formula of **X**?

(1)

- A** C_7H_{16}
- B** $C_6H_{12}O$
- C** $C_6H_{14}N$
- D** $C_5H_8O_2$

(ii) The infrared spectrum of **X** contains major absorption wavenumber ranges at $3300\text{--}2500\text{ cm}^{-1}$, $1725\text{--}1700\text{ cm}^{-1}$ and $1669\text{--}1645\text{ cm}^{-1}$.

Identify the two functional groups in **X**.

(2)

(iii) **X** has an unbranched carbon chain and does **not** exhibit geometric isomerism.

Draw the **skeletal formula** of **X**.

(1)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 6 2 6 6 9 A 0 1 1 3 2



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

A large rectangular area with horizontal dotted lines for writing, spanning most of the page width and height.


(Total for Question 8 = 10 marks)



P 6 2 6 6 9 A 0 1 1 3 2




9 This question is about the effect of temperature on the rate of decomposition of nitrogen(V) oxide.



(a) The diagram shows the Maxwell-Boltzmann distribution of molecular energies for nitrogen(V) oxide at a temperature T_1 .

E_a is the activation energy of this reaction.



(i) Give the label for the vertical axis. (1)

(ii) Draw a second curve on the same set of axes for the same gas at a **lower** temperature, T_2 . (2)

(iii) Explain, in terms of collisions and energy, why lowering the temperature decreases the rate of reaction. (2)

.....

.....

.....

.....

.....

.....

.....

.....

(iv) A catalyst is added to the gas.

Label the diagram above with the symbol E_{cat} to show a possible activation energy for the reaction in the presence of a catalyst. (1)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





- (b) The rate constant for the decomposition of nitrogen(V) oxide was determined at two temperatures.

Temperature / K	Rate constant / s ⁻¹
328	1.50 × 10 ⁻³
338	4.87 × 10 ⁻³

Calculate the activation energy for this reaction.

Include units and give your answer to an appropriate number of significant figures.

You should **not** attempt to use any graphical method to answer this question.

The Arrhenius equation relating two rate constants, k_1 and k_2 , at two different temperatures, T_1 and T_2 , can be expressed as

$$\ln\left(\frac{k_2}{k_1}\right) = -\frac{E_a}{R}\left(\frac{1}{T_2} - \frac{1}{T_1}\right) \quad (5)$$

(Total for Question 9 = 11 marks)

TOTAL FOR PAPER = 90 MARKS





DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



P 6 2 6 6 9 A 0 2 9 3 2



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



P 6 2 6 6 9 A 0 2 9 3 2

The Periodic Table of Elements

1 2 3 4 5 6 7 0 (8)
(18)

1.0	H
	hydrogen
1	

Key

relative atomic mass
atomic symbol
name
atomic (proton) number

(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
6.9	9.0	45.0	47.9	50.9	52.0	54.9	55.8	58.9	58.7	63.5	65.4	10.8	12.0	14.0	16.0	19.0	4.0
Li lithium 3	Be beryllium 4	Sc scandium 21	Ti titanium 22	V vanadium 23	Cr chromium 24	Mn manganese 25	Fe iron 26	Co cobalt 27	Ni nickel 28	Cu copper 29	Zn zinc 30	B boron 5	C carbon 6	N nitrogen 7	O oxygen 8	F fluorine 9	He helium 2
23.0	24.3	88.9	91.2	92.9	95.9	[98]	101.1	102.9	106.4	107.9	112.4	27.0	28.1	31.0	32.1	35.5	39.9
Na sodium 11	Mg magnesium 12	Y yttrium 39	Zr zirconium 40	Nb niobium 41	Mo molybdenum 42	Tc technetium 43	Ru ruthenium 44	Rh rhodium 45	Pd palladium 46	Ag silver 47	Cd cadmium 48	Al aluminium 13	Si silicon 14	P phosphorus 15	S sulfur 16	Cl chlorine 17	Ar argon 18
39.1	40.1	138.9	178.5	180.9	183.8	186.2	190.2	192.2	195.1	197.0	200.6	69.7	72.6	74.9	79.0	79.9	83.8
K potassium 19	Ca calcium 20	La* lanthanum 57	Hf hafnium 72	Ta tantalum 73	W tungsten 74	Re rhenium 75	Os osmium 76	Ir iridium 77	Pt platinum 78	Au gold 79	Hg mercury 80	Ga gallium 31	Ge germanium 32	As arsenic 33	Se selenium 34	Br bromine 35	Kr krypton 36
85.5	87.6	[227]	178.5	180.9	183.8	186.2	190.2	192.2	195.1	197.0	200.6	114.8	118.7	121.8	127.6	126.9	131.3
Rb rubidium 37	Sr strontium 38	Ac* actinium 89	Hf hafnium 72	Ta tantalum 73	W tungsten 74	Re rhenium 75	Os osmium 76	Ir iridium 77	Pt platinum 78	Au gold 79	Hg mercury 80	In indium 49	Sn tin 50	Sb antimony 51	Te tellurium 52	I iodine 53	Xe xenon 54
132.9	137.3	[227]	178.5	180.9	183.8	186.2	190.2	192.2	195.1	197.0	200.6	204.4	207.2	209.0	[209]	[210]	[222]
Cs caesium 55	Ba barium 56	Ra radium 88	Hf hafnium 72	Ta tantalum 73	W tungsten 74	Re rhenium 75	Os osmium 76	Ir iridium 77	Pt platinum 78	Au gold 79	Hg mercury 80	Tl thallium 81	Pb lead 82	Bi bismuth 83	Po polonium 84	At astatine 85	Rn radon 86
[223]	[226]	[227]	[261]	[262]	[266]	[264]	[277]	[268]	[271]	[272]							
Fr francium 87	Ra radium 88	Ac* actinium 89	Rf rutherfordium 104	Db dubnium 105	Sg seaborgium 106	Bh bohrium 107	Hs hassium 108	Mt meitnerium 109	Ds darmstadtium 110	Rg roentgenium 111							

Elements with atomic numbers 112-116 have been reported but not fully authenticated

140	141	144	150	152	157	159	163	165	167	169	173	175
Ce cerium 58	Pr praseodymium 59	Nd neodymium 60	Sm samarium 62	Eu europium 63	Gd gadolinium 64	Tb terbium 65	Dy dysprosium 66	Ho holmium 67	Er erbium 68	Tm thulium 69	Yb ytterbium 70	Lu lutetium 71
232	[231]	238	[242]	[243]	[247]	[245]	[251]	[254]	[253]	[256]	[254]	[257]
Th thorium 90	Pa protactinium 91	U uranium 92	Pu plutonium 94	Am americium 95	Cm curium 96	Bk berkelium 97	Cf californium 98	Es einsteinium 99	Fm fermium 100	Md mendelevium 101	No nobelium 102	Lr lawrencium 103

* Lanthanide series

* Actinide series

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

