



Please check the examination details below before entering your candidate information

Candidate surname

Other names

Pearson
Edexcel GCE

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--

Thursday 23 May 2019

Morning (Time: 1 hour 30 minutes)

Paper Reference **8CH0/02**

Chemistry
Advanced Subsidiary
Paper 2: Core Organic and Physical Chemistry

Candidates must have: **Scientific calculator**
Data Booklet
Ruler

Total Marks

Instructions

- Use **black** ink or **black** ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
- *there may be more space than you need.*

Information

- The total mark for this paper is 80.
- The marks for **each** question are shown in brackets
- *use this as a guide as to how much time to spend on each question.*
- For the question marked with an **asterisk** (*), marks will be awarded for your ability to structure your answer logically showing the points that you make are related or follow on from each other where appropriate.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Check your answers if you have time at the end.

Turn over ►

P55611A

©2019 Pearson Education Ltd.

2/2/1/1/1/



www.ibworldwideacademy.com



Pearson



Answer ALL questions.

Some questions must be answered with a cross in a box ☒.
If you change your mind about an answer, put a line through the box ☒
and then mark your new answer with a cross ☒.

1 Some alkanes are used as fuels for transport. Crude oil, which is a mixture of hydrocarbons, is the major source of these alkanes. Crude oil is processed by fractional distillation, cracking and reforming.

(a) Fractional distillation produces fractions which contain molecules of a similar boiling temperature. Molecules containing six carbon atoms are found in both the petrol fraction and the higher boiling kerosene fraction.

(i) Identify, by name or structural formula, the unbranched alkane with six carbon atoms found in kerosene.

(1)

.....

(ii) Explain why **isomers** of the alkane in (a)(i) have lower boiling temperatures and so are found in the petrol fraction.

(3)

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 5 5 6 1 1 A 0 3 2 8



(b) During cracking, carbon to carbon bonds are broken.

Explain the **two** major reasons for cracking hydrocarbons.

(3)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Which of the following changes to the proportions of organic molecules within a fraction results from the reforming process?

(1)

	Proportion of branched chain alkanes	Proportion of cyclic hydrocarbons
<input type="checkbox"/> A	decrease	decrease
<input type="checkbox"/> B	decrease	increase
<input type="checkbox"/> C	increase	decrease
<input type="checkbox"/> D	increase	increase

(Total for Question 1 = 8 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 5 5 6 1 1 A 0 3 2 8



2 This question is about organic compounds.

(a) Organic compounds can be grouped together in homologous series.

(i) Describe **two** characteristics of a homologous series.

(2)

.....

.....

.....

.....

.....

.....

(ii) Name the homologous series to which propene belongs.

(1)

.....

(b) Propene can be converted into a mixture of 1-chloropropane and 2-chloropropane, in which 2-chloropropane is the major product.

(i) Give the reagent required for this reaction.

(1)

.....

(ii) What is the type and mechanism of the reaction in (b)(i)?

(1)

- A electrophilic addition
- B nucleophilic addition
- C electrophilic substitution
- D nucleophilic substitution

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 5 5 6 1 1 A 0 3 2 8



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(c) 1-chloropropane and 2-chloropropane can be converted into compounds containing the nitrile functional group.

(i) Under appropriate conditions, 1-chloropropane can be converted into butanenitrile, $\text{CH}_3\text{CH}_2\text{CH}_2\text{CN}$.

Which is the reagent for this conversion?

(1)

- A ammonia
- B nitric acid
- C potassium cyanide
- D silver nitrate

(ii) Under appropriate conditions, 2-chloropropane can be converted into a structural isomer of butanenitrile.

State what is meant by the term 'structural isomer'.

(2)

.....

.....

.....

.....

(iii) Give the displayed formula **and** systematic name of the isomer of butanenitrile formed in (c)(ii). You must show **all** the bonds.

(2)

Displayed formula

Name

.....

(Total for Question 2 = 10 marks)



P 5 5 6 1 1 A 0 3 2 8



3 This question is about enthalpy changes.

(a) (i) State what is meant by the term 'standard enthalpy change of combustion'.

(2)

.....

.....

.....

.....


(ii) Write the equation, including state symbols, for the reaction that occurs when the standard enthalpy change of combustion of octane, $C_8H_{18}(l)$, is measured.

(2)

(iii) The standard enthalpy change of combustion of octane is $-5470 \text{ kJ mol}^{-1}$.

Complete the reaction profile diagram for the combustion of octane. Include labels showing the standard enthalpy change of combustion, ΔH^\ominus , and the activation energy, E_a .

(2)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 5 5 6 1 1 A 0 3 2 8



- (b) Enthalpy changes of reactions which cannot be measured directly can be calculated using standard enthalpy changes of combustion. The table shows some of these values.

Substance	$\Delta_c H^\circ / \text{kJ mol}^{-1}$
C(s)	-394
H ₂ (g)	-286
CH ₄ (g)	-890

Complete the Hess cycle and use it to calculate the standard enthalpy change for the following reaction.

(4)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 5 5 6 1 1 A 0 3 2 8





DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

- (c) The equations for the combination of gaseous atoms of carbon and hydrogen to form methane, CH₄, and propane, C₃H₈, are



Calculate:

- (i) the mean bond enthalpy of a C--H bond. (1)

- (ii) the mean bond enthalpy of a C--C bond. (2)

(Total for Question 3 = 13 marks)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



P 5 5 6 1 1 A 0 9 2 8



4 Methanol is manufactured from a mixture of carbon monoxide and hydrogen.



(a) Give **two** characteristics of all reactions at equilibrium.

(2)

.....

.....

.....

.....

.....

(b) (i) How does the equilibrium yield of methanol change if the temperature is increased at constant pressure or the pressure increased at constant temperature?

(1)

	Equilibrium yield when temperature is increased	Equilibrium yield when pressure is increased
<input type="checkbox"/> A	decrease	decrease
<input type="checkbox"/> B	decrease	increase
<input type="checkbox"/> C	increase	decrease
<input type="checkbox"/> D	Increase	increase

(ii) Explain your answer to (b)(i).

(2)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





(c) Explain why, in the industrial process involving this reaction, a catalyst is used.

(2)

.....

.....

.....

.....

.....

.....

.....

(Total for Question 4 = 7 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA




P 5 5 6 1 1 A 0 1 1 2 8



5 Bromobutanes react with hot ethanolic potassium hydroxide solution to produce gaseous butenes.



Apparatus



Procedure

- 0.0080 mol of liquid 1-bromobutane was injected into a round bottom flask containing hot ethanolic potassium hydroxide.
- After the reaction, the syringe was sealed using a clamp.
- The syringe was then removed from the apparatus and allowed to cool to room temperature (298 K).

Result

The final volume of but-1-ene collected was 22.0 cm³.

(a) State the purpose of the condenser.

(1)

.....

.....

.....

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(b) Describe a chemical test on the gas in the syringe to identify its functional group.
Include the expected result.

(2)

.....

.....

.....

.....

.....

.....

.....

(c) Calculate the percentage of 1-bromobutane which was converted to but-1-ene.
[Molar volume of a gas at r.t.p. = $24\,000\text{ cm}^3\text{ mol}^{-1}$]

(2)

(d) Before cooling, the volume of but-1-ene in the gas syringe was 24.0 cm^3 .
Calculate the temperature of the gas in the syringe before it cooled.

[Assume no loss from the gas syringe during cooling, and a constant pressure]

(2)



P 5 5 6 1 1 A 0 1 1 2 8



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(e) (i) Another compound formed from 1-bromobutane under these conditions is butan-1-ol. Identify the type of reaction taking place to form butan-1-ol. (1)

(ii) The functional group in butan-1-ol can be confirmed using a single chemical test. What is the single chemical test and expected observation? (1)

	Chemical test reagent	Observation
<input type="checkbox"/> A	sodium carbonate solution	effervescence
<input type="checkbox"/> B	aqueous silver nitrate	cream precipitate
<input type="checkbox"/> C	Fehling's solution	red precipitate
<input type="checkbox"/> D	phosphorus(V) chloride	steamy fumes

(iii) Draw the mechanism for the reaction of 1-bromobutane with hydroxide ions to form butan-1-ol.

Include curly arrows, and any appropriate lone pairs and dipoles. (3)





(f) Alkene molecules are formed by elimination from 2-bromobutane.

How many isomeric alkene products will be formed in this reaction?

(1)

- A 1
- B 2
- C 3
- D 4

(Total for Question 5 = 13 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 5 5 6 1 1 A 0 1 1 2 8



6 Wine and gin are aqueous solutions of ethanol with traces of other organic compounds which give these drinks their characteristic flavours and aromas.

(a) When a bottle of wine is opened, oxidation of the ethanol in the wine produces ethanoic acid.

An experiment was carried out to determine the percentage of the ethanol that had been oxidised.

- A bottle of white wine, with an ethanol concentration of 2.50 mol dm^{-3} , was opened and left to stand at room temperature for three weeks.
- A 25.0 cm^3 sample of the wine was transferred to a conical flask and phenolphthalein indicator added.
- Aqueous sodium hydroxide of concentration $0.235 \text{ mol dm}^{-3}$ was added from a burette until the colour of the indicator permanently changed.
- The titration was repeated and the titre values, in cm^3 , were 27.90, 26.75 and 26.85.

The equation for the neutralisation reaction is



(i) Name the piece of apparatus used to measure 25.0 cm^3 of wine.

(1)

(ii) To improve the accuracy, the burette should be rinsed.

State what should be used.

(1)

(iii) What is the colour change at the end-point?

(1)

- A from orange to yellow
- B from red to orange
- C from colourless to pink
- D from pink to colourless

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





(iv) State what is meant by the term 'concordant results' as applied to a titration experiment.

(1)

.....

.....

(v) Calculate the percentage of ethanol that has oxidised, given that one mole of ethanol forms one mole of ethanoic acid.

Give your answer to an appropriate number of significant figures.

(5)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 5 5 6 1 1 A 0 1 1 2 8



(vi) Deduce why this method would **not** be effective for the analysis of the acid content of a red wine.

(1)

.....

.....

(b) The ethanol content of alcoholic drinks is usually measured as the percentage of alcohol by volume (ABV).

$$\text{ABV} = \frac{\text{Volume of ethanol in a solution}}{\text{Total volume of the solution}} \times 100$$

The ABV values of four different brands of gin are shown.

Brand of gin	ABV (%)
A	40
B	42
C	44
D	46

A sample of one of these gins was found to contain an ethanol concentration of 7.50 mol dm^{-3} .

By calculating the percentage of ethanol by volume (ABV) of this sample, deduce the brand of this gin.

[Assume the density of ethanol, $\text{C}_2\text{H}_5\text{OH} = 0.79 \text{ g cm}^{-3}$]

(3)

(Total for Question 6 = 13 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE





7 This question is about compounds with the molecular formula C_4H_8O .

(a) What is the percentage by mass of each element in C_4H_8O ?

(1)

	Percentage carbon	Percentage hydrogen	Percentage oxygen
<input type="checkbox"/> A	66.67	11.11	22.22
<input type="checkbox"/> B	60.00	20.00	20.00
<input type="checkbox"/> C	41.38	3.45	55.17
<input type="checkbox"/> D	30.77	61.54	7.69

*(b) Three different compounds, **A**, **B** and **C**, have the molecular formula C_4H_8O .

Information about these three compounds includes:

- all three compounds have infrared absorptions at about 3500 cm^{-1}
- the infrared spectra of **A** and **B** each contain a peak at about 1650 cm^{-1} , while that of **C** does not
- only **A** has a branched carbon chain
- **B** is the *E*-isomer of a pair of stereoisomers.

Deduce a possible **displayed** formula for each of the compounds **A**, **B** and **C**.

You must use **all** the information and the Data Booklet to fully justify each of your structures.

(6)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 5 5 6 1 1 A 0 2 1 2 8



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

A large rectangular area with horizontal dotted lines for writing, spanning most of the page width and height.

(Total for Question 7 = 7 marks)



P 5 5 6 1 1 A 0 2 1 2 8



8 Compound **X** reacts slowly with water according to the following equation.



The reaction is catalysed by hydrogen ions and eventually goes to completion.

Compound **X** was added to water and the concentration of compound **Y** determined at various times at a constant temperature.

The results of the experiment are shown.

Time / s	Concentration of Y / mol dm ⁻³
0	0.000
25	0.002
40	0.005
50	0.010
65	0.020
75	0.030
85	0.040
100	0.050
115	0.060
140	0.070
200	0.080
260	0.085
380	0.090

- (a) (i) Complete the graph of concentration against time by adding the six missing points.
Draw a line to pass through **all** the points.

(2)

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





Graph of concentration of **Y** against time



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 5 5 6 1 1 A 0 2 1 2 8



(ii) Describe how you would find a numerical value for the initial rate of reaction and for the maximum rate of reaction in this experiment from the graph. No actual calculations are required.

(4)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(b) For many reactions, the values of the initial rate and the maximum rate are the same.

Explain why the values of the two reaction rates obtained in this experiment are different from each other.

(2)

.....

.....

.....

.....

.....

.....

(c) Give a reason why the measurement of the initial rate of reaction is likely to be less accurate than the measurement of the maximum rate.

(1)

.....

.....

.....

(Total for Question 8 = 9 marks)

TOTAL FOR PAPER = 80 MARKS



P 5 5 6 1 1 A 0 2 1 2 8

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



BLANK PAGE



P 5 5 6 1 1 A 0 2 6 2 8

BLANK PAGE



The Periodic Table of Elements

1 2 3 4 5 6 7 0 (8) (18)

6.9	Li	lithium	3	(1)
9.0	Be	beryllium	4	(2)
23.0	Na	sodium	11	(3)
24.3	Mg	magnesium	12	(3)
39.1	K	potassium	19	(4)
40.1	Ca	calcium	20	(4)
45.0	Sc	scandium	21	(5)
47.9	Ti	titanium	22	(5)
50.9	V	vanadium	23	(6)
52.0	Cr	chromium	24	(6)
54.9	Mn	manganese	25	(7)
55.8	Fe	iron	26	(8)
58.9	Co	cobalt	27	(9)
58.7	Ni	nickel	28	(10)
63.5	Cu	copper	29	(11)
65.4	Zn	zinc	30	(12)
69.7	Ga	gallium	31	(13)
72.6	Ge	germanium	32	(14)
74.9	As	arsenic	33	(15)
79.0	Se	selenium	34	(16)
79.9	Br	bromine	35	(17)
83.8	Kr	krypton	36	(18)
131.3	Xe	xenon	54	(18)
126.9	I	iodine	53	(18)
127.6	Te	tellurium	52	(18)
121.8	Sb	antimony	51	(18)
118.7	Sn	tin	50	(18)
114.8	In	indium	49	(18)
112.4	Cd	cadmium	48	(18)
107.9	Ag	silver	47	(18)
106.4	Pd	palladium	46	(18)
102.9	Rh	rhodium	45	(18)
101.1	Ru	ruthenium	44	(18)
[98]	Tc	technetium	43	(18)
95.9	Mo	molybdenum	42	(18)
92.9	Nb	niobium	41	(18)
91.2	Zr	zirconium	40	(18)
88.9	Y	yttrium	39	(18)
87.6	Sr	strontium	38	(18)
85.5	Rb	rubidium	37	(18)
87.3	Ba	barium	56	(18)
137.3	La*	lanthanum	57	(18)
138.9	Ce	cerium	58	(18)
132.9	Pr	praseodymium	59	(18)
137.3	Nd	neodymium	60	(18)
178.5	Pm	promethium	61	(18)
178.5	Sm	samarium	62	(18)
180.9	Eu	europium	63	(18)
183.8	Gd	gadolinium	64	(18)
186.2	Tb	terbium	65	(18)
187.7	Dy	dysprosium	66	(18)
190.2	Ho	holmium	67	(18)
192.2	Er	erbium	68	(18)
195.1	Tm	thulium	69	(18)
197.0	Yb	ytterbium	70	(18)
197.0	Lu	lutetium	71	(18)
200.6	Hf	hafnium	72	(18)
200.6	Ta	tantalum	73	(18)
204.4	W	tungsten	74	(18)
204.4	Re	rhenium	75	(18)
207.2	Os	osmium	76	(18)
207.2	Ir	iridium	77	(18)
209.0	Pt	platinum	78	(18)
209.0	Au	gold	79	(18)
209.0	Hg	mercury	80	(18)
209.0	Tl	thallium	81	(18)
209.0	Pb	lead	82	(18)
209.0	Bi	bismuth	83	(18)
209.0	Po	polonium	84	(18)
210	At	astatine	85	(18)
222	Rn	radon	86	(18)

140	Ce	cerium	58	(18)
141	Pr	praseodymium	59	(18)
144	Nd	neodymium	60	(18)
147	Pm	promethium	61	(18)
150	Sm	samarium	62	(18)
152	Eu	europium	63	(18)
157	Gd	gadolinium	64	(18)
159	Tb	terbium	65	(18)
163	Dy	dysprosium	66	(18)
165	Ho	holmium	67	(18)
167	Er	erbium	68	(18)
169	Tm	thulium	69	(18)
173	Yb	ytterbium	70	(18)
175	Lu	lutetium	71	(18)
140	Th	thorium	90	(18)
232	Pa	protactinium	91	(18)
238	U	uranium	92	(18)
237	Np	neptunium	93	(18)
242	Pu	plutonium	94	(18)
243	Am	americium	95	(18)
247	Cm	curium	96	(18)
245	Bk	berkelium	97	(18)
251	Cf	californium	98	(18)
254	Es	einsteinium	99	(18)
253	Fm	fermium	100	(18)
256	Md	mendelevium	101	(18)
254	No	nobelium	102	(18)
257	Lr	lawrencium	103	(18)

* Lanthanide series
* Actinide series

Elements with atomic numbers 112-116 have been reported but not fully authenticated

relative atomic mass
atomic symbol
name
atomic (proton) number

1.0
H
hydrogen
1

